arXiv:1612.00967v1 [cs.IT] 3 Dec 2016

# Two new families of two-weight codes

Minjia Shi

*Key Laboratory of Intelligent Computing & Signal Processing

Ministry of Education, Anhui University No. 3 Feixi Road

Hefei Anhui Province 230039, P. R. China;
National Mobile Communications Research Laboratory

Southeast University, 210096, Nanjing, P. R. China;

School of Mathematical Sciences

Anhui University, Hefei, 230601, P. R. China
smjwcl.good@163.com

Yue Guan

School of Mathematical Sciences,
Anhui University, Hefei, 230601, P. R. China

Patrick Solé

CNRS/ LAGA
University Paris 8, 93 526 Saint-Denis, France
patrick.sole@telecom-paristech.fr

**Abstract**

We construct two new infinite families of trace codes of dimension $2m$, over the ring $\mathbb{F}_p + u\mathbb{F}_p$, when $p$ is an odd prime. They have the algebraic structure of abelian codes. Their Lee weight distribution is computed by using Gauss sums. By Gray mapping, we obtain two infinite families of linear $p$-ary codes of respective lengths $(p^m-1)^2$ and $2(p^m-1)^2$. When $m$ is singly-even, the first family gives five-weight codes. When $m$ is odd, and $p \equiv 3 \pmod 4$, the first family yields $p$-ary two-weight codes, which are shown to be optimal by application of the Griesmer bound. The second family consists of two-weight codes that are shown to be optimal, by the Griesmer bound, whenever $p = 3$ and $m \geq 3$, or $p \geq 5$ and $m \geq 4$. Applications to secret sharing schemes are given.

**Keywords:** Two-weight codes, Gauss sums, Griesmer bound, secret sharing schemes

*This research is supported by National Natural Science Foundation of China (61672036), the Open Research Fund of National Mobile Communications Research Laboratory, Southeast University (2015D11), Technology Foundation for Selected Overseas Chinese Scholar, Ministry of Personnel of China (05015133) and Key projects of support program for outstanding young talents in Colleges and Universities (gxyqZD2016008).

# 1 Introduction

Two-weight codes over fields have been studied since the 1970s due to their connections to strongly regular graphs, finite geometries and difference sets [5]. However, most constructions, have used cyclic codes [3, 4]. In the present paper, we use trace codes over a semi-local ring which is a quadratic extension of a finite field, and obtain codes over a finite field by Gray mapping. This is part of a general research program where a variety of few weight codes are obtained by varying the base ring and the defining set [14, 15, 16]. Here, we keep the base ring fixed, but we consider two families depending on two different defining sets. These codes are visibly abelian, but possibly not cyclic. The field image of the first family has two or five weights, depending on the choice of parameters. The second family only contains two-weight codes. Note that abelian codes over rings have been studied already in [11]. The alphabet ring we consider here is $\mathbb{F}_p + u\mathbb{F}_p$, with $u^2 = u$. It is a semi-local ring, abstractly isomorphic to $\mathbb{F}_p \times \mathbb{F}_p$. It is an odd characteristic analogue of $\mathbb{F}_{2^r} + v\mathbb{F}_{2^r}$, with $v^2 = v$. The latter ring has been employed recently to construct convolutional codes over fields [9]. Some bounds on codes over $\mathbb{F}_p + u\mathbb{F}_p$, with $u^2 = u$ can be found in [10].

The defining set of our abelian code is not a cyclic group, but it is an abelian group. The defining set of the first family is related to quadratic residues in an extension of degree $m$ of $\mathbb{F}_p$, which makes quadratic Gauss sums appear naturally in the weight distribution analysis, and requires $p$ to be an odd prime. When $m$ is odd, and $p \equiv 3 \pmod 4$, we obtain an infinite family of linear $p$-ary two-weight codes, which are shown to be optimal by application of the Griesmer bound. The codes in the second family are also shown to be optimal by the same technique up to finitely many exceptions. We show that, both in the five-weight and in the two-weight cases, the first family has a very nice support inclusion structure which makes it suitable for use in a Massey secret sharing scheme [7, 8, 18]. Indeed, we can show that all nonzero codewords are minimal for the poset of codewords ordered by support inclusion. A similar result holds for the second family.

The paper is organized as follows. Section 2 collects the notions and notations needed in the rest of the article. Section 3 shows that the trace codes are abelian. Section 4 recalls and reproves some results on Gaussian periods. Section 5 computes the weight distribution of our codes, building on the character sum evaluation of the preceding section. Section 6 discusses the optimality of the $p$-image from trace codes over $R$. Section 7 determines the minimum distance of the dual codes. Section 8 determines the support structure of the $p$-ary image and describes an application to secret sharing schemes.

# 2 Definitions and notations

## 2.1 Rings

Consider the ring $R = \mathbb{F}_p + u\mathbb{F}_p$ where $u^2 = u$ and $p$ is a odd prime. It is semi-local with maximal ideals $(u)$ and $(u-1)$. For any integer $m \geq 1$, we construct an extension of degree $m$ as $\mathcal{R} = \mathbb{F}_{p^m} + u\mathbb{F}_{p^m}$ with again $u^2 = u$. There is a Frobenius operator $F$ which maps $a + ub$ onto $a^p + ub^p$. The *Trace function,* denoted by $Tr$ is then defined as

$Tr = \sum_{j=0}^{m-1} F^j$. It is then immediate from these definitions that $Tr(a+ub) = tr(a)+utr(b)$, for $a, b \in \mathbb{F}_{p^m}$. Here $tr()$ denotes the absolute trace of $\mathbb{F}_{p^m}$, given by

$$tr(z) = z + z^p + \cdots + z^{p^{m-1}}, z \in \mathbb{F}_{p^m}.$$

The ring $\mathcal{R}$ is semi-local with maximal ideals $(u)$ and $(u-1)$. The Chinese Remainder Theorem shows then that it is $u\mathbb{F}_{p^m}+(1-u)\mathbb{F}_{p^m}$, isomorphic to the product ring $\mathbb{F}_{p^m} \times \mathbb{F}_{p^m}$. Similarly, the group of units $\mathcal{R}^*$ is $u\mathbb{F}_{p^m}^* + (1-u)\mathbb{F}_{p^m}^*$, isomorphic to $\mathbb{F}_{p^m}^* \times \mathbb{F}_{p^m}^*$. The *squares* and the *non-squares* of $\mathbb{F}_{p^m}$ are denoted by $\mathcal{Q}$ and $\mathcal{N}$, respectively. We write $L = u\mathcal{Q} + (1-u)\mathbb{F}_{p^m}^*$ and, for simplicity, we let $L' = \mathcal{R}^*$. Thus $L$ is a subgroup of $\mathcal{R}^*$ of index 2.

### 2.2 Gray map

The *Gray* map $\phi$ from $R$ to $\mathbb{F}_p^2$ is defined by $\phi(a + ub) = (-b, 2a + b)$ for $a, b \in \mathbb{F}_p$. It is a one to one map from $R$ to $\mathbb{F}_p^2$, which extends naturally into a map from $R^n$ to $\mathbb{F}_p^{2n}$. The *Lee weight* is defined as the Hamming weight of the Gray image $w_L(a + ub) = w_H(-b) + w_H(2a+b)$, for $a, b \in \mathbb{F}_p^n$. The *Lee distance* of $x, y \in R^n$ is defined as $w_L(x-y)$. Thus the Gray map is a linear isometry from $(R^n, d_L)$ to $(\mathbb{F}_p^{2n}, d_H)$. For convenience, we write $N = 2n$ in the rest of the paper.

### 2.3 Codes

A **linear code** $C$ over $R$ of length $n$ is an $R$-submodule of $R^n$. If $x = (x_1, x_2, \cdots, x_n)$ and $y = (y_1, y_2, \cdots, y_n)$ are two elements of $R^n$, their standard inner product is defined by $\langle x, y\rangle = \sum_{i=1}^{n} x_i y_i$, where the operation is performed in $R$. The **dual code** of $C$ is denoted by $C^\perp$ and defined as $C^\perp = \{y \in R^n | \langle x, y\rangle = 0, \forall x \in C\}$. By definition, $C^\perp$ is also a linear code over $R$. Given a finite abelian group $G$, a code over $R$ is said to be **abelian** [2], if it is an ideal of the group ring $R[G]$. Recall that the ring $R[G]$ is defined on functions from $G$ to $R$ with pointwise addition as addition, and convolution product as multiplication. Concretely, it is the set of all formal sums $f = \sum_{h\in G} f_h X^h$, with addition and multiplication defined as follows. If $f, g$ are in $R[G]$, we write

$$f + g = \sum_{g\in G} (f_h + g_h) X^h,$$

and

$$fg = \sum_{h\in G} \Big( \sum_{r+s=h} f_r g_s \Big) X^h.$$

In other words, the coordinates of $C$ are indexed by elements of $G$ and $G$ acts regularly on this set. For more details on abelian codes see [17]. In the special case when $G$ is cyclic, the code is a cyclic code in the usual sense [12].

## 3 Symmetry

For $a \in \mathcal{R}$ define the vector $ev(a)$ by the following evaluation map $ev(a) = (Tr(ax))_{x \in L}$. Define the code $C(m,p)$ by the formula $C(m,p) = \{ev(a) | a \in \mathcal{R}\}$. Thus $C(m,p)$ is a code of length $|L| = \frac{(p^m-1)^2}{2}$, and size $|R|^m$ over $R$. Similarly define the vector $ev'(a)$ by the following evaluation map $ev'(a) = (Tr(ax))_{x \in L'}$, and the code $C'(m,p)$ by the formula $C'(m,p) = \{ev'(a) | a \in \mathcal{R}\}$. Thus $C'(m,p)$ is a code of length $|L'| = (p^m - 1)^2$ and size $|R|^m$ over $R$.

**Proposition 1.** *The group $L$ (resp. $L'$ ) acts regularly on the coordinates of $C(m,p)$ (resp. $C'(m,p)$ ).*

*Proof.* For any $w, v \in L$ the change of variables $x \mapsto (v/w)x$ maps $w$ to $v$. This transformation defines thus a transitive action of $L$ on itself. Given an ordered pair $(w,v)$ this transformation is unique, hence the action is regular. A similar argument holds for $C'(m,p)$ and $L'$. □

The code $C(m,p)$ is thus an *abelian code* with respect to the group $\mathcal{R}^*$. In other words, it is an ideal of the group ring $R[\mathcal{R}^*]$. As observed in the previous section $\mathcal{R}^*$ is not a cyclic group, hence $C(m,p)$ may be not cyclic.

## 4 Character sums

Let $\chi$ denote an arbitrary multiplicative character of $\mathbb{F}_q$. Assume $q$ is odd. Denoted by $\eta$ the quadratic multiplicative character is defined by $\eta(x) = 1$, if $x$ is a square and $\eta(x) = -1$, if not. Let $\psi$ denote the standard canonical additive character of $\mathbb{F}_q$. The classical **Gauss sum** can be defined as $G(\chi) = \sum_{x \in \mathbb{F}_q^*} \psi(x)\chi(x)$.

We define the following character sums

$$\overline{Q} = \sum_{x \in \mathcal{Q}} \psi(x), \ \overline{N} = \sum_{x \in \mathcal{N}} \psi(x).$$

By orthogonality of characters [12, Lemma 9, p. 143], it is easy to see that $\overline{Q} + \overline{N} = -1$. Noting that the characteristic function of $\mathcal{Q}$ is $\frac{1+\eta}{2}$, we get then

$$\overline{Q} = \frac{G(\eta) - 1}{2}, \ \overline{N} = \frac{-G(\eta) - 1}{2}.$$

It is well known [6] that if $q = p^m$, the quadratic Gauss sums can be evaluated as

$$G(\eta) = (-1)^{m-1}\sqrt{q}, \ p \equiv 1 \pmod 4, \quad (1)$$
$$G(\eta) = (-1)^{m-1} i^m \sqrt{q}, \ p \equiv 3 \pmod 4. \quad (2)$$

Particularly, if $m$ is singly-even, these formulas can be simplified to $G(\eta) = \epsilon(p)\sqrt{q}$, with $\epsilon(p) = (-1)^{\frac{(p+1)}{2}}$, yielding

$$\overline{Q} = \frac{\epsilon(p)\sqrt{q} - 1}{2}, \overline{N} = -\frac{\epsilon(p)\sqrt{q} + 1}{2}.$$

In fact $\overline{Q}$ and $\overline{N}$ are examples of *Gaussian periods*, and these relations could have been deduced from [6, Lemma 11].

# 5 Weight distributions of trace codes

We begin with a correlation lemma. Let $\omega = \exp(\frac{2\pi i}{p})$. If $y = (y_1, y_2, \cdots, y_N) \in \mathbb{F}_p^N$, let $\Theta(y) = \sum_{j=1}^{N} \omega^{y_j}$. For simplicity, we let $\theta(a) = \Theta(\phi(ev(a)))$. By linearity of the Gray map, and of the evaluation map, we see that $\theta(sa) = \Theta(\phi(ev(sa)))$, for any $s \in \mathbb{F}_p^*$.

**Lemma 2.** *[15] For all $y = (y_1, y_2, \cdots, y_N) \in \mathbb{F}_p^N$, we have $\sum_{s=1}^{p-1} \Theta(sy) = (p-1)N - pw_H(y)$.*

## 5.1 The case of $L = \mathcal{Q} \times \mathbb{F}_{p^m}^*$

### 5.1.1 $m$ is singly-even

**Theorem 3.** *Assume $m$ is singly-even. For $a \in \mathcal{R}$, the Lee weight of codewords of $C(m,p)$ is given below.*

*(a) If $a = 0$, then $w_L(ev(a)) = 0$;*

*(b) If $a = u\alpha$, $\alpha \in \mathbb{F}_{p^m}^*$, then if*

$\alpha \in \mathcal{Q}$ *then* $w_L(ev(a)) = (p-1)\big(p^{2m-1} - p^{m-1} - \epsilon(p)p^{3m/2-1} + \epsilon(p)p^{m/2-1}\big)$,
$\alpha \in \mathcal{N}$ *then* $w_L(ev(a)) = (p-1)\big(p^{2m-1} - p^{m-1} + \epsilon(p)p^{3m/2-1} - \epsilon(p)p^{m/2-1}\big)$;

*(c) If $a = (1-u)\beta$, $\beta \in \mathbb{F}_{p^m}^*$, then $w_L(ev(a)) = (p-1)(p^{2m-1} - p^{m-1})$;*

*(d) If $a = u\alpha + (1-u)\beta \in \mathcal{R}^*$, then if*

$\alpha \in \mathcal{Q}$ *then* $w_L(ev(a)) = (p-1)\big(p^{2m-1} - 2p^{m-1} + \epsilon(p)p^{m/2-1}\big)$,
$\alpha \in \mathcal{N}$ *then* $w_L(ev(a)) = (p-1)\big(p^{2m-1} - 2p^{m-1} - \epsilon(p)p^{m/2-1}\big)$.

*Proof.* (a) If $a = 0$, then $Tr(ax) = 0$. So $w_L(ev(a)) = 0$.

(b) If $a = u\alpha$, $x = ut + (1-u)t'$ with $\alpha \in \mathbb{F}_{p^m}^*$, then $ax = u\alpha t$, and $Tr(ax) = Tr(u\alpha t) = utr(\alpha t)$. Taking Gray map yields $\phi(ev(a)) = (-tr(\alpha t), tr(\alpha t))_{t,t'}$. Taking character sums

$$\begin{aligned}
\theta(a) &= \sum_{t\in\mathcal{Q}} \sum_{t'\in\mathbb{F}_{p^m}^*} \omega^{-tr(\alpha t)} + \sum_{t\in\mathcal{Q}} \sum_{t'\in\mathbb{F}_{p^m}^*} \omega^{tr(\alpha t)} \\
&= 2\sum_{t\in\mathcal{Q}} \sum_{t'\in\mathbb{F}_{p^m}^*} \omega^{-tr(\alpha t)} \\
&= 2(p^m - 1)\sum_{t\in\mathcal{Q}} \omega^{tr(\alpha t)}.
\end{aligned}$$

Replaced $\alpha t$ by $t$, it is easy to check that the last character sum is $\overline{Q}$ or $\overline{N}$ depending on $\alpha \in \mathcal{Q}$ or $\alpha \in \mathcal{N}$. Since $m$ is even, $s \in \mathbb{F}_p^*$ is a square in $\mathbb{F}_{p^m}$. Thus $\theta(sa) = \theta(a)$, for any $s \in \mathbb{F}_p^*$. The statement follows from Lemma 2. Thus $w_L(ev(a)) = \frac{p-1}{p}(N - 2(p^m-1)\overline{Q})$, or $w_L(ev(a)) = \frac{p-1}{p}(N - 2(p^m-1)\overline{N})$, according to the value of $\eta(\alpha)$.

(c) If $a = (1-u)\beta$, $x = ut + (1-u)t'$ with $\beta \in \mathbb{F}_{p^m}^*$, then $ax = \beta t' - u\beta t'$, and $Tr(ax) = tr(\beta t') - utr(\beta t')$. Taking Gray map yields $\phi(ev(a)) = (tr(\beta t'), tr(\beta t'))_{t,t'}$. Taking character sums $\theta(a) = 2\sum\limits_{t\in\mathcal{Q}}\sum\limits_{t'\in\mathbb{F}_{p^m}^*}\omega^{tr(\beta t')} = 1 - p^m$. Thus $w_L(ev(a)) = (p-1)(p^{2m-1} - p^{m-1})$.

(d) Let $a = u\alpha + (1-u)\beta \in R^*$, $x = ut + (1-u)t'$. So $Tr(ax) = tr(\beta t') + utr(\alpha t - \beta t')$. Taking Gray map yields $\phi(ev(a)) = (-tr(\alpha t - \beta t'), tr(\alpha t + \beta t'))_{t,t'}$. Taking character sums $\theta(a) = \sum\limits_{t\in\mathcal{Q}}\omega^{-tr(\alpha t)}\sum\limits_{t'\in\mathbb{F}_{p^m}^*}\omega^{tr(\beta t')} + \sum\limits_{t\in\mathcal{Q}}\omega^{tr(\alpha t)}\sum\limits_{t'\in\mathbb{F}_{p^m}^*}\omega^{tr(\beta t')} = -2\sum\limits_{t\in\mathcal{Q}}\omega^{tr(\alpha t)}$. By a change of variable $t = \alpha t$, we see that the last character sum is $\overline{Q}$ or $\overline{N}$ depending on $\alpha \in \mathcal{Q}$ or $\alpha \in \mathcal{N}$. Thus $w_L(ev(a)) = \frac{p-1}{p}(N + 2\overline{Q})$, or $w_L(ev(a)) = \frac{p-1}{p}(N + 2\overline{N})$, considering the value of $\eta(\alpha)$.

□

Therefore, we have constructed a $p$-ary code of length $N = (p^m-1)^2$, dimension $2m$, with five weights. The weight distribution is given in Table I.

Table I. weight distribution of $C(m,p)$ in Theorem 3

| Weight | Frequency |
|---|---|
| 0 | 1 |
| $(p-1)(p^{m-1} - p^{m/2-1})(p^m-1)$ | $\frac{p^m-1}{2}$ |
| $(p-1)(p^{2m-1} - 2p^{m-1} - p^{m/2-1})$ | $\frac{(p^m-1)^2}{2}$ |
| $(p-1)(p^{2m-1} - 2p^{m-1} + p^{m/2-1})$ | $\frac{(p^m-1)^2}{2}$ |
| $(p-1)(p^{2m-1} - p^{m-1})$ | $p^m - 1$ |
| $(p-1)(p^{m-1} + p^{m/2-1})(p^m-1)$ | $\frac{p^m-1}{2}$ |

(Note that taking $\epsilon(p) = 1$, or $-1$, leads to the same values.)

### 5.1.2 $m$ is odd and $p \equiv 3 \pmod 4$

Note that in that case by (2) in Section 4 we see that $G(\eta)$ is imaginary. This implies that $\Re(\overline{Q}) = \Re(\overline{N}) = -\frac{1}{2}$. We need first to refine the following correlation lemma.

**Lemma 4.** *[15] If $p \equiv 3 \pmod 4$, then we have $\sum\limits_{s=1}^{p-1}\theta(sa) = (p-1)\Re(\theta(a))$.*

**Theorem 5.** *Assume $m$ is odd and $p \equiv 3 \pmod 4$. For $a \in \mathcal{R}$, the Lee weight of codewords of $C(m,p)$ is given below.*

*(a) If $a = 0$, then $w_L(ev(a)) = 0$;*

*(b) If $a = u\beta$, then $w_L(ev(a)) = (p-1)(p^{2m-1} - p^{m-1})$;*

*(c) If $a = (1-u)\beta$, then $w_L(ev(a)) = (p-1)(p^{2m-1} - p^{m-1})$;*

*(d) If $a \in \mathcal{R}^*$, then $w_L(ev(a)) = (p-1)(p^{2m-1} - 2p^{m-1})$.*

*Proof.* The proof of the case (a) is like that of Theorem 3. The case (b) follows from Lemma 4 applied to the correlation lemma. Thus $\Re(\theta(a)) = 1 - p^m$, and $pw_L(ev(a)) = (p-1)(N - \Re(\theta(a)))$, yielding $w_L(ev(a)) = (p-1)(p^{2m-1} - p^{m-1})$. The result follows. The proof of case (c) is the same as that of case (b). In the case (d), $\Re(\theta(a)) = 1$, then $w_L(ev(a)) = (p-1)(p^{2m-1} - 2p^{m-1})$. □

Thus we obtain a family of $p$-ary two-weight codes of parameters $[p^{2m} - 2p^m + 1, 2m]$, with weight distribution as given in Table II. The parameters are different from those in [4], [14], [15] and [16].

Table II. weight distribution of $C(m,p)$ in Theorem 5

| Weight | Frequency |
|---|---|
| 0 | 1 |
| $(p-1)(p^{2m-1} - 2p^{m-1})$ | $(p^m-1)^2$ |
| $(p-1)(p^{2m-1} - p^{m-1})$ | $2(p^m-1)$ |

### 5.2 The case of $L' = \mathbb{F}^*_{p^m} \times \mathbb{F}^*_{p^m}$

**Theorem 6.** *For $a \in \mathcal{R}$, the Lee weight of codewords of $C'(m,p)$ is*

*(a) If $a = 0$, then $w_L(ev'(a)) = 0$;*

*(b) If $a = u\alpha$, $\alpha \in \mathbb{F}^*_{p^m}$, then $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - p^{m-1})$;*

*(c) If $a = (1-u)\beta$, $\beta \in \mathbb{F}^*_{p^m}$, then $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - p^{m-1})$;*

*(d) If $a \in \mathcal{R}^*$, then $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - 2p^{m-1})$.*

*Proof.* (a) If $a = 0$, then $Tr(ax) = 0$. So $w_L(ev'(a)) = 0$.

(b) If $a = u\alpha$, $x = ut + (1-u)t'$ with $\alpha \in \mathbb{F}^*_{p^m}$, then $ax = u\alpha t$, and $Tr(ax) = Tr(u\alpha t) = utr(\alpha t)$. Taking Gray map yields $\phi(ev'(a)) = (-tr(\alpha t), tr(\alpha t))_{t,t'}$. Taking character sums $\theta(a) = \sum_{t \in \mathbb{F}^*_{p^m}} \sum_{t' \in \mathbb{F}^*_{p^m}} \omega^{-tr(\alpha t)} + \sum_{t \in \mathbb{F}^*_{p^m}} \sum_{t' \in \mathbb{F}^*_{p^m}} \omega^{tr(\alpha t)} = 2 \sum_{t \in \mathbb{F}^*_{p^m}} \sum_{t' \in \mathbb{F}^*_{p^m}} \omega^{-tr(\alpha t)} = -2(p^m - 1)$. Thus $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - p^{m-1})$.

(c) If $a = (1-u)\beta$, $x = ut + (1-u)t'$ with $\beta \in \mathbb{F}^*_{p^m}$, then $ax = \beta t' - u\beta t'$, and $Tr(ax) = tr(\beta t') - utr(\beta t')$. Taking Gray map yields $\phi(ev'(a)) = (tr(\beta t'), tr(\beta t'))_{t,t'}$. Taking character sums $\theta(a) = 2 \sum_{t \in \mathbb{F}^*_{p^m}} \sum_{t' \in \mathbb{F}^*_{p^m}} \omega^{tr(\beta t')} = 2 - 2p^m$. Thus $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - p^{m-1})$.

(d) Let $a = u\alpha + (1-u)\beta)$, $x = ut + (1-u)t'$. So $ax = \beta t' + u(\alpha t - \beta t')$, and $Tr(ax) = tr(\beta t') + utr(\alpha t - \beta t')$. Taking Gray map yields $\phi(ev'(a)) = \big( - tr(\alpha t - \beta t'), tr(\alpha t + \beta t')\big)_{t,t'}$. Taking character sums $\theta(a) = \sum_{t \in \mathbb{F}_{p^m}^*} \omega^{-tr(\alpha t)} \sum_{t' \in \mathbb{F}_{p^m}^*} \omega^{tr(\beta t')} + \sum_{t \in \mathbb{F}_{p^m}^*} \omega^{tr(\alpha t)} \sum_{t' \in \mathbb{F}_{p^m}^*} \omega^{tr(\beta t')} = 2$. Thus $w_L(ev'(a)) = \frac{p-1}{p}(N' - 2)$, or $w_L(ev'(a)) = 2(p-1)(p^{2m-1} - 2p^{m-1})$.

□

Thus we have constructed a $p$-ary code of length $N' = 2(p^m - 1)^2$, dimension $2m$, with two weights. The weight distribution is given in Table III. Note that the parameters are different from those in [4], [14], [15] and [16].

Table III. weight distribution of $C'(m,p)$ in Theorem 6

| Weight | Frequency |
|---|---|
| 0 | 1 |
| $2(p-1)(p^{2m-1} - 2p^{m-1})$ | $p^{2m} - 2p^m + 1$ |
| $2(p-1)(p^{2m-1} - p^{m-1})$ | $2p^m - 2$ |

## 6 Optimality of the $p$-ary image

We now study their optimality. Recall the $p$-ary version of the Griesmer bound. If $[N, K, d]$ are the parameters of a linear $p$-ary code, then

$$\sum_{j=0}^{K-1} \left\lceil \frac{d}{p^j} \right\rceil \leq N.$$

### 6.1 $L = u\mathcal{Q} + (1-u)\mathbb{F}_{p^m}^*$, $m$ is odd and $p \equiv 3 \pmod 4$

**Theorem 7.** *Assume $m$ is odd and $m \geq 3$, and $p \equiv 3 \pmod 4$. The code $\phi(C(m,p))$ is optimal.*

*Proof.* In our situation $N = p^{2m} - 2p^m + 1$, $K = 2m$, $d = (p-1)(p^{2m-1} - 2p^{m-1})$. We claim that $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil > N$, violating the Griesmer bound. The ceiling function takes three values depending on $j$.

- $0 \leq j \leq m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = p^{2m-j} - 2p^{m-j} - p^{2m-j-1} + 2p^{m-j-1} + 1$;
- $j = m \Rightarrow \lceil \frac{d+1}{p^j} \rceil = p^m - p^{m-1} - 1$;
- $m < j \leq 2m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = p^{2m-j} - p^{2m-j-1}$.

Thus $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil = p^{2m} - 2p^m + m$. Note that $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil - N = m - 1 > 0$. □

**Example 8.** Let $p = 3$ and $m = 3$, we obtain a ternary code of parameters $[676, 6, 450]$. The weights of this code are 450 and 468 with frequencies 676 and 52, respectively.

### 6.2 $L' = \mathcal{R}^*$, $m$ is an arbitrary integer and $p$ is odd prime

**Theorem 9.** *Assume $m \geq 3$ and $p = 3$ or $m \geq 4$ and the odd prime $p \geq 5$. The code $\phi(C'(m,p))$ is optimal.*

*Proof.* In the situation $N' = 2(p^{2m} - 2p^m + 1)$, $K = 2m$, $d = 2(p-1)(p^{2m-1} - 2p^{m-1})$. We claim that $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil > N'$, violating the Griesmer bound. The ceiling function takes the following values depending on the position of $j$.

(a) when $p = 3$, the ceiling function takes three values depending on $j$.

- $0 \leq j \leq m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^{2m-j} - 4p^{m-j} - 2p^{2m-j-1} + 4p^{m-j-1} + 1$;
- $j = m \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^m - 2p^{m-1} - 2$;
- $m < j \leq 2m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^{2m-j} - 2p^{2m-j-1}$.

Thus $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil = 2p^{2m} - 4p^m + m$. Note that $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil - N' = m - 2 > 0$.

(b) when $p \geq 5$ and $p$ is odd prime, the ceiling function takes three values depending on $j$.

- $0 \leq j \leq m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^{2m-j} - 4p^{m-j} - 2p^{2m-j-1} + 4p^{m-j-1} + 1$;
- $j = m \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^m - 2p^{m-1} - 3$;
- $m < j \leq 2m-1 \Rightarrow \lceil \frac{d+1}{p^j} \rceil = 2p^{2m-j} - 2p^{2m-j-1}$.

Thus $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil = 2p^{2m} - 4p^m + m - 1$. Note that $\sum_{j=0}^{K-1} \lceil \frac{d+1}{p^j} \rceil - N' = m - 3 > 0$.

Hence the theorem is proved. □

**Example 10.** Let $p = 3$ and $m = 3$, we obtain a ternary code of parameters $[1352, 6, 900]$. The weights of this code are 900 and 936 with frequencies 676 and 52, respectively.

## 7 The minimum distance of the dual code

We compute the dual distance of $\phi(C(m,p))$(resp. $\phi(C'(m,p))$). The proof of the following lemma is similar to that in [14].

**Lemma 11.** *If for all $a \in \mathcal{R}$, we have that $Tr(ax) = 0$, then $x = 0$.*

**Theorem 12.** *For all odd primes $p$ and all $m \geq 2$, the dual Lee distance $d'$ of $C(m,p)$ is 2.*

*Proof.* First, we check that $d' \geq 2$, by showing that $C(m,p)^{\perp}$, does not contain a word of Lee weight one. If it does, let us assume first that it has value $\gamma \neq 0$ at some $x \in L$. This implies that $\forall a \in \mathcal{R}, \gamma Tr(ax) = 0$, or, $Tr(a\gamma x) = 0$, and by Lemma 11 $\gamma x = 0$. Contradiction with $\gamma \neq 0$. If that word takes the value $\gamma(1-2u)$ at some $x \in L$, then writing $x = ut + (1-u)t'$ and $a = u\alpha + (1-u)\beta$, with $\alpha, \beta$ in $\mathbb{F}_{p^m}$, with $t \in \mathcal{Q}, t' \in \mathbb{F}_{p^m}^*$ yields, after reduction modulo $M$ the equation valid $\forall \alpha, \beta \in \mathbb{F}_{p^m}$, $\gamma tr(\beta t') = 0$ and $tr(\gamma\alpha t) = 0$, and we conclude, by the nondegenerate character of $tr()$, that $\gamma t = \gamma t' = 0$. Contradiction with $x \neq 0$.

Next, we show that $d' < 3$. If not, we can apply the sphere-packing bound to $\phi(C(m,p))^{\perp}$, to obtain $p^{2m} \geq 1 + N(p-1) = 1 + (p^{2m} - 2p^m + 1)(p-1)$, or, after expansion $2p^{2m} + 2p^{m+1} \geq p + p^m(p^{m+1} + 2)$. Dropping the $p$ in the RHS, and dividing both sides by $p^m$, we find that this inequality would imply $p^m < f(p)$, with $f(x) = \frac{2x-2}{x-2} = 2 + \frac{2}{x-2}$. But the function $f$ is decreasing for $x \geq 3$, and $f(3) = 4$, while $p^m \geq p^2 \geq 9$. Contradiction. □

We now consider codes from the second family.

**Theorem 13.** *For all odd prime $p$ and all $m \geq 2$, the dual Lee distance $d'$ of $C'(m,p)$ is 2.*

*Proof.* First, we check that $d' \geq 2$. The proof of $d' \geq 2$ is like that in Theorem 12. Next, we show that $d' < 3$. If not, we can apply the sphere-packing bound to $\phi(C'(m,p)^{\perp})$, to obtain

$$p^{2m} \geq 1 + N'(p-1) = 1 + 2(p^{2m} - 2p^m + 1)(p-1).$$

It is easy to check that $1 + N'(p-1) > 1 + N'$ where $p$ is an odd prime. Then we check $p^{2m} > 1 + N'$. We obtain the inequality $(p^m - 3)(p^m - 1) < 0$ after calculation which is contradict $p^m \geq 3$. Hence $p^{2m} \leq 1 + N'(p-1) = 1 + 2(p^{2m} - 2p^m + 1)(p-1)$. □

## 8 Applications to secret sharing schemes

The support $s(x)$ of a vector $x$ in $\mathbb{F}_p^N$ is defined as the set of indices where it is nonzero. We say that a vector $x$ covers a vector $y$ if $s(x)$ contains $s(y)$. A *minimal codeword* of a linear code $C$ is a nonzero codeword that does not cover any other nonzero codeword. In general, it is a tough task to determine the minimal codewords of a given linear code. However, there is a numerical condition, derived in [1], bearing on the weights of the code, that is easy to check.

**Lemma 14.** *(Ashikmin-Barg) Let $w_0$ and $w_\infty$ denote the minimum and maximum nonzero weights, respectively. If $\frac{w_0}{w_\infty} > \frac{p-1}{p}$, then every nonzero codeword of $C$ is minimal.*

We can infer from there the support structure for the codes of this paper. For the first family, in the five-weight case, we obtain the result below.

**Proposition 15.** *All the nonzero codewords of $\phi(C(m,p))$ are minimal when $m(\geq 6)$ is singly-even.*

*Proof.* By the preceding lemma with $w_0 = (p-1)(p^{m-1} - p^{m/2-1})(p^m - 1)$ and $w_\infty = (p-1)(p^{m-1} + p^{m/2-1})(p^m - 1)$ in Table I. We end up with the condition $p^{m/2} + 1 > 2p$, which is true for $m \geq 6$ and $p$ is an odd prime. □

The analogue for the two-weight code family is as follows.

**Proposition 16.** *All the nonzero codewords of $\phi(C(m,p))$, for $m$ odd $\geq 3$, and $p \equiv 3 \pmod 4$, are minimal.*

*Proof.* By the preceding lemma with $w_0 = (p-1)(p^{2m-1} - 2p^{m-1})$ and $w_\infty = (p-1)(p^{2m-1} - p^{m-1})$ in Table II. Rewriting the inequality of the lemma as $pw_0 > (p-1)w_\infty$, and dividing both sides by $\frac{p^m}{p-1}$, we obtain $p(p^m - 2) > (p-1)(p^m - 1)$, or $p^m - p - 1 > 0$, which is true for $m \geq 3$. □

**Proposition 17.** *All the nonzero codewords of $\phi(C'(m,p))$, for all $m \geq 2$, and odd prime $p$ are minimal.*

*Proof.* By the preceding lemma with $w_0 = 2(p-1)(p^{2m-1} - 2p^{m-1})$ and $w_\infty = 2(p-1)(p^{2m-1} - p^{m-1})$ in Table III. Simplify the inequality $pw_0 > (p-1)w_\infty$ as $p^m - p - 1 > 0$, which is true for $m \geq 2$. □

Massey's scheme [18], is a secret sharing scheme, based on Coding theory. In the special case when all nonzero codewords are minimal, it was shown in [7] that there is the following alternative, depending on $d'$:

- If $d' \geq 3$, then the SSS is *"democratic"*: every user belongs to the same number of coalitions.
- If $d' = 2$, then there are users who belong to every coalition: the *"dictators"*.

By Propositions 15, 16, 17, and Theorems 12, 13, we see that three Secret Sharing Schemes built on $\phi(C(m,p))$ or $\phi(C'(m,p))$ are dictatorial.

## 9 Conclusion Remarks

In the present paper, we have studied two infinite families of trace codes defined over a finite ring. Because the defining sets of these codes have the structure of abelian multiplicative groups, they inherit the structure of abelian codes. It is an open problem to determine if they are cyclic codes or not. More importantly, it is worthwhile to study other defining sets that are subgroups of the group of units of $\mathcal{R}$. In particular, it would be interesting to replace our Gaussian periods $\overline{Q}, \overline{N}$ by other character sums that are amenable to exact evaluation, in the vein of the sums which appear in the study of irreducible cyclic codes [6, 13]. This would lead to other enumerative results of codes with a few weights.

Compared to the codes we constructed by similar techniques in [14], [15] and [16], the obtained codes in this paper have different weight distributions. They are also different from the weight distributions in the classical families in [4]. Hence, the $p$-ary linear codes constructed from trace codes over rings in this paper are new.

## References


[1] A. Ashikmin, A. Barg, "Minimal vectors in linear codes", *IEEE Trans. Inform. Theory*, vol. 44, no. 5, pp. 2010–2017, Sep. 1998.

[2] D. Berman, "Semisimple cyclic and Abelian codes. II", *Cybernetics & Systems Analysis*, vol. 3, no. 3, pp. 17–23, 1967.

[3] A.E. Brouwer, W.H. Haemers, "Spectra of Graphs", *Springer*, 2012.

[4] R. Calderbank, W.M. Kantor, "The geometry of two-weight codes", *Bull. London Math. Soc.*, vol. 18, no. 2, pp. 97-122, 1986.

[5] P. Delsarte, "Weights of linear codes and strongly regular normed spaces", *Disc. Math.*, vol. 3, no. 1-3, pp. 47–64, 1972.

[6] C. Ding, J. Yang, "Hamming weights in irreducible cyclic codes", *Disc. Math.*, vol. 313, issue 4, pp. 434–446, Feb. 2013.

[7] C. Ding, J. Yuan, "Covering and secret sharing with linear codes", *Springer LNCS*, vol. 2731, pp. 11-25, Jun. 2003.

[8] K.L. Ding, and C. Ding, "A Class of Two-Weight and Three-Weight Codes and Their Applications in Secret Sharing", *IEEE Trans. Inform. Theory*, vol. 61, issue 11, pp. 5835–5842, Nov. 2015.

[9] J.M. Lampos, V. Sison, "Systematic Unit-Memory Binary Convolutional Codes from Linear Block Codes over $\mathbb{F}_{2^r} + v\mathbb{F}_{2^r}$", *WASET*, issue 71, pp. 384-387, Nov. 2012.

[10] J.M. Lampos, V. Sison, "Bounds on (..) linear block codes over finite semi-local Frobenius ring $\mathbb{F}_{p^r} + v\mathbb{F}_{p^r}$", Southeast-Asian J. of Sc., Vol. 1, No 1 (2012), pp.76–98.

[11] S. Ling, P. Solé, "Duadic codes over $\mathbb{F}_2 + u\mathbb{F}_2$", *AAECC*, vol. 12, no. 5, pp. 365-379, Oct. 2001.

[12] F.J. MacWilliams, N.J.A. Sloane, "The theory of error-correcting codes", *North-Holl Math. Libr*, vol. 16, 1977.

[13] R.J. McEliece, H.Jr. Rumsey, "Euler products, cyclotomy, and coding", *J. Number Theory*, vol. 4, no. 3, pp. 302–311, Jun. 1972.

[14] M.J. Shi, Y. Liu, P. Solé, "Optimal two weight codes from trace codes over $\mathbb{F}_2+u\mathbb{F}_2$", *IEEE Commun. Letters*, http://ieeexplore.ieee.org/document/7582413/.

[15] M.J. Shi, Y. Liu, P. Solé, Optimal two weight codes from trace codes over a non-chain ring, *Discrete Applied Mathematics*, Accepted for publication.

[16] M.J. Shi, R.S. Wu, Y. Liu, P. Solé, "two and three weight codes over $\mathbb{F}_p + u\mathbb{F}_p$", *Cryptogr. Commun.*, doi:10.1007/s12095-016-0206-5, Sep. 2016.

[17] H.N. Ward, "Quadratic residue codes an divisibility, in Handbook of Coding Theory, H. Ward and V. Pless eds", *North Holland*, 827–870, 1998.

[18] J. Yuan, C. Ding, "Secret sharing schemes from three classes of linear codes", *IEEE Trans. Inform. Theory*, vol. 52, no.1, pp. 206-212, Jan. 2006.